

WITH GRAPH PAPER

केन्द्रीय माध्यमिक शिक्षा बोर्ड, दिल्ली
आचार्य कृष्ण सचिवालय परीक्षा (कक्षा दसवीं)
परिभाषी प्रश्न-पत्र के अनुसार एवं

Mathematics

041

I

Schedule 18.03.2015

English

Code Number

65/2/c

Set Number

1 2 3 4

Yes / No

40

B D H S C A

Yes / No

40

Each letter be written in one box and one box be left blank between each part of the answer. In case Candidates Name exceeds 24 letters, write first 24 letters.

341-0852 4414974

Section - 1

Q: 20

$$(1+x^2) \frac{dy}{dx} = e^{\tan^{-1}x} - y$$

$$(1+x^2) \frac{dy}{dx} + y = e^{\tan^{-1}x}$$

$$\frac{dy}{dx} + \frac{y}{1+x^2} = \frac{e^{\tan^{-1}x}}{1+x^2}$$

comparing it with

$$\frac{dy}{dx} + Py = Q \rightarrow \text{linear differential equation}$$

$$P = \frac{1}{1+x^2}$$

$$I.O.F = e^{\int P dx}$$

$$= e^{\int \frac{1}{1+x^2} dx}$$

$$I.O.F = e^{\tan^{-1}x}$$

$$\left\{ \int \frac{1}{x^2+1} dx = \tan^{-1}x \right\}$$

the diff equation becomes

$$y \text{ I.O.F} = \int Q \times \text{I.O.F} \, dx + c$$

$$y \times e^{\tan^{-1}x} = \int \frac{e^{\tan^{-1}x}}{1+x^2} \times e^{\tan^{-1}x} \, dx + c$$

$$y \times e^{\tan^{-1}x} = \int \frac{e^{(m+1)\tan^{-1}x}}{1+x^2} \, dx + c$$

but $\tan^{-1}x = t$

$$\frac{1}{1+x^2} \, dx = dt$$

$$y e^{\tan^{-1}x} = \int e^{(m+1)t} \times dt + c$$

$$y e^{\tan^{-1}x} = \frac{e^{(m+1)t}}{(m+1)} + c$$

$$\text{after } y e^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{m+1} + c$$

when $x=0$ $y=1$

$$ye^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{(m+1)} + c$$

when $x=0$ $y=1$

$$1 \times e^{\tan^{-1}0} = \frac{e^{(m+1)\tan^{-1}0}}{(m+1)} + c$$

$$1 \times e^0 = \frac{e^{(m+1)0}}{m+1} + c \quad e^0 = 1$$

$$1 = \frac{1}{m+1} + c$$

$$1 - \frac{1}{m+1} = c$$

$$\frac{m+1-1}{m+1} = c$$

$$c = \frac{m}{m+1}$$

the equation is

$$\left[ye^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{(1+m)} + \frac{m}{m+1} \right]$$

Q: 21.

$$f(x) = \sin^2 x - \cos x \quad x \in [0, \pi]$$

$$f'(x) = 2 \sin x \cos x - (-\sin x)$$

$$f'(x) = 2 \sin x \cos x + \sin x$$

$$\text{put } f'(x) = 0$$

$$\text{then } (2 \sin x \cos x + \sin x) = 0$$

$$\sin x (2 \cos x + 1) = 0$$

$$\sin x = 0 \quad \text{or} \quad \cos x = -\frac{1}{2}$$

$$x = 0, \pi \quad \text{or} \quad x = \frac{2\pi}{3}$$

$$\text{also } f(0) = [\sin(0)]^2 - \cos 0$$

$$= 0 - 1$$

$$= -1$$

$$f(\pi) = \sin^2 \pi - \cos \pi$$

$$= 0 - (-1)$$

$$= 1$$

$$f\left(\frac{2\pi}{3}\right) = \sin^2 \frac{2\pi}{3} - \cos \frac{2\pi}{3}$$

$$= \left(\frac{\sqrt{3}}{2}\right)^2 - \left(-\frac{1}{2}\right)$$

$$= \frac{3}{4} + \frac{1}{2} = \frac{10}{8} = \frac{5}{4}$$

both the extreme value are automatically included

$$f(0) = -1$$

$$f(1) = 1$$

$$f\left(\frac{20}{3}\right) = \frac{5}{4}$$

So Absolute maxima is $\frac{5}{4}$ at $x = \frac{20}{3}$

Absolute minima is -1 at $x = 0$

Q: 22

$$R = 1 + j + k + \lambda(1 - j + k)$$

$$R = 4j + 2k + \mu(2i - j + 3k)$$

These line are coplanar if they are parallel or they are intersecting

But these lines are not parallel.

So they are coplanar if shortest distance between them is zero.

$$a_1 = (1, 1, 1)$$

$$a_2 = (0, 4, 2)$$

$$b_1 = \langle 1, -1, 1 \rangle$$

$$b_2 = \langle 2, -1, 3 \rangle$$

$$(a_2 - a_1) = (-1, 3, 1)$$

$$\text{Shortest distance} = \frac{|(a_2 - a_1) \cdot (b_1 \times b_2)|}{|b_1 \times b_2|}$$

Let us find $|(a_2 - a_1) \cdot (b_1 \times b_2)|$

$$(a_2 - a_1) \cdot (b_1 \times b_2) =$$

$$\begin{vmatrix} -1 & 3 & 1 \\ 1 & -1 & 1 \\ 2 & -1 & 3 \end{vmatrix}$$

(expanding by 1st row)

$$= -1(-3+1) - 3(3-2) + 1(-1+2)$$

$$= -1(-2) - 3(1) + 1$$

$$= 2 + 1 - 3$$

$$= 0$$

So shortest distance between the lines is zero
So they are coplanar.

Let $\langle a, b, c \rangle$ be the direction ratio of normal to the plane.

So the dot product of direction ratio of normal to plane and the direction ratio of line is 0.

So $a - b + c = 0$

$2a - b + 3c = 0$

<u>a</u>	<u>b</u>	<u>c</u>
-1	1	-1
-1	3	-1

$$\frac{a}{-3+1} = \frac{b}{2-3} = \frac{c}{-1+2}$$

So $\langle a, b, c \rangle = \langle -2, -1, 1 \rangle$

So passing point of line is also the passing point of plane

So passing point = $(1, 1, 1)$

the equation becomes

$$-2(x-1) - 1(y-1) + 1(z-1) = 0$$

$$-2x + 2 - y + 1 + z - 1 = 0$$

$$\boxed{-2x - y + z + 2 = 0}$$

equation of plane.

Q: 23

$$f: W \rightarrow W$$

$$f(n) = \begin{cases} n-1 & \text{if } n \text{ is odd} \\ n+1 & \text{if } n \text{ is even} \end{cases}$$

prove that it is one-one.

Let us suppose that $n_1, n_2 \in W$

$$n_1 \neq n_2 \text{ but } f(n_1) = f(n_2)$$

(i) 1st case : if n_1 & n_2 are odd

$$f(n_1) = f(n_2)$$

$$n_1 - 1 = n_2 - 1$$

which is contradiction.

(ii) 2nd case : n_1, n_2 are even

$$f(n_1) = f(n_2)$$

$$n_1 + \cancel{1} = n_2 + \cancel{1}$$

$$n_1 = n_2$$

which is contradiction.

(iii) 3rd case if n_1 is even & n_2 is odd

$$f(n_1) = f(n_2)$$

$$n_1 + 1 = n_2 - 1$$

$$n_1 + 2 = n_2$$

$$\begin{array}{ccc} \downarrow & & \downarrow \\ \text{even} & & \text{odd} \end{array}$$

if we add 2 in an even no. then we get an even no. but n_2 is odd which is again contradiction

So from these 3 point we see that

$$\Rightarrow f(n_1) = f(n_2) \text{ only if } n_1 = n_2$$

$\Rightarrow f$ is one-one.

To prove that f is onto:

let $y \in \mathbb{N}$

and y is even

then $y+1$ is odd and it belongs in vehicle no.

$$f(y+1) = y+1-1$$

$$= y$$

So for every $y \in \mathbb{N}$ which is even there exist a preimage $y+1$ which is odd and $(y+1) \in \mathbb{N}$

let $y \in \mathbb{N}$

y is odd

then $y-1$ is even and $y-1 \in \mathbb{N}$

$$f(y-1) = y-1+1$$

$$= y$$

So for every $y \in \mathbb{N}$ which is odd there exist a preimage $y-1 \in \mathbb{N}$ which is even.

So for all $y \in \mathbb{N}$ there exist preimage in vehicle no. So f is onto.

f is one-one and onto so it is invertible

To find $f^{-1}(x)$

$$y = x - 1 \quad \text{if } x \text{ is odd}$$

$$y + 1 = x$$

$$\text{so } x = y + 1 \quad \text{if } y \text{ is even}$$

$$f^{-1}(x) = x + 1 \quad \text{if } x \text{ is even}$$

$$y = x + 1 \quad \text{if } x \text{ is even}$$

$$y - 1 = x \quad \text{if } y \text{ is odd here}$$


$$y - 1 = x$$

$$\text{so } f^{-1}(x) = x - 1 \quad \text{if } x \text{ is odd}$$

$$\text{so } f^{-1}(x) = \begin{cases} x - 1 & \text{if } x \text{ is odd} \\ x + 1 & \text{if } x \text{ is even} \end{cases}$$

$$[x \in W]$$

$$\text{so } [f^{-1} = f]$$

240

intersection point :

$$y = \sqrt{5-x^2}$$

$$y = x-1$$

$$\text{so } \sqrt{5-x^2} = (x-1)$$

$$5-x^2 = x^2+1-2x$$

$$2x^2-2x-4=0$$

$$2x^2-x-2=0$$

$$x^2-2x+x-2=0$$

$$x(x-2)+1(x-2)=0$$

$$\text{so } x=2 \text{ or } x=-1$$

$$x=2$$

rejected as $x \geq 1$

$$y = \sqrt{5-x^2}$$

$$y = -x+1$$

$$y \quad x < 1$$

$$5-x^2 = x^2 + 1 - 2x$$

$$\text{So } x = 2, 0 - 1$$

$$x = 2$$

$$[x = -1]$$

Rejected

Required Area

$$(A) = 2 \int_{-1}^2 \sqrt{5-x^2} dx - \left[\int_{-1}^2 (-x+1) dx + \int_1^2 (x-1) dx \right]$$

$$A = \frac{x\sqrt{5-x^2}}{2} + \frac{5}{2} \sin^{-1} \frac{x}{\sqrt{5}} \Big|_{-1}^2 - \left[\frac{-x^2+x}{2} \Big|_{-1}^2 + \frac{x^2-x}{2} \Big|_1^2 \right]$$

$$A = 1 \times 1 + \frac{5}{2} \sin^{-1} \frac{2}{\sqrt{5}} - \left(\frac{-1 \times 2}{2} + \frac{5}{2} \sin^{-1} \left(\frac{-1}{\sqrt{5}} \right) \right) - \left[\frac{-1}{2} [2-1] + 2 \right. \\ \left. + \frac{1}{2} [4-1] - 1 \right]$$

$$A = \frac{1+5 \sin^{-1} 2}{2} + \frac{1+5 \sin^{-1} 1}{2} - \left[0 + 2 + \frac{3}{2} - 1 \right]$$

$$A = \frac{-2+5 \left[\sin^{-1} 2 + \sin^{-1} 1 \right] - 5}{2} = \frac{5}{2}$$

$$A = \frac{5 \times \frac{\pi}{2} + 2 - 5}{2}$$

$$A = \frac{5\pi}{4} - \frac{1}{2} \text{ 89 units}$$

$$\left\{ \begin{aligned} & \frac{\sin^{-1} 2}{\sqrt{5}} + \frac{\sin^{-1} 1}{\sqrt{5}} \\ &= \frac{\sin^{-1} 2 + \sin^{-1} 1}{\sqrt{5}} \\ &= \frac{\pi}{2} \end{aligned} \right\}$$

$\int_0^1 dx$

250

Let 6 positive integers = (1, 2, 3, 4, 5, 6)

Random Variable = X = larger of 2 numbers.

$$x \left| \begin{array}{c} 2 \\ 1 \end{array} \right.$$

2

$$\{4-1\}-1\}$$

sample space $S = \{$

$$\begin{aligned} & (1,2) (1,3) (1,4) (1,5) (1,6) \\ & (2,1) (3,1) (4,1) (5,1) (6,1) \\ & (2,3) (2,4) (2,5) (2,6) (3,2) (4,2) (5,2) \\ & (6,2) (3,4) (3,5) (3,6) (4,3) (5,3) (6,3) \\ & (4,5) (4,6) (5,4) (6,4) (5,6) (6,5) \end{aligned}$$

X	P(X)	$X \cdot P_i$	$X^2 \cdot P_i$	
2	$\frac{2}{30} = \frac{1}{15}$	$\frac{2}{15}$	$\frac{4}{15}$	
3	$\frac{4}{30} = \frac{2}{15}$	$\frac{6}{15}$	$\frac{18}{15}$	
4	$\frac{6}{30} = \frac{3}{15}$	$\frac{12}{15}$	$\frac{48}{15}$	
5	$\frac{8}{30} = \frac{4}{15}$	$\frac{20}{15}$	$\frac{100}{15}$	
6	$\frac{10}{30} = \frac{5}{15}$	$\frac{30}{15}$	$\frac{180}{15}$	
				Range 2+6

$$\sum X_i P_i = 70 \quad \sum X_i^2 P_i = 350$$

$$\text{Mean} = \frac{\sum X_i P_i}{n} = \frac{70}{15} = 4.66$$

$$\text{Variance} = \frac{\sum X_i^2 P_i - (\mu)^2}{n}$$

$$= \frac{350 - 70 \times 70}{15}$$

$$= \frac{350 - 4900}{15}$$

$$s^2 = \frac{70}{15} \left[5 - \frac{70}{15} \right]$$

$$= \frac{70}{15} \left[\frac{75-70}{15} \right]$$

$$= \frac{70 \times 5}{315}$$

$$= \frac{14}{9}$$

$$s^2 = 1.55$$


$$\text{Variance} = 1.55$$

Section - B

Q:19



$$x^x + x^y + y^x = a^b$$



$$\frac{d}{dx} [P + Q + R] = 0$$

$$\left[\frac{dx}{dx} + \frac{dy}{dx} + \frac{dx}{dx} = 0 \right]$$

$$P = x^x$$

$$\log P = x \log x$$

$$\frac{1}{P} \frac{dP}{dx} = x \left[\frac{1}{x} \right] + \log x$$

$$\frac{dP}{dx} = x^x (1 + \log x) \quad - (1)$$

$$Q = x^y$$

$$\log Q = y \log x$$

$$\frac{1}{Q} \frac{dQ}{dx} = \left(\frac{y}{x} + \log x \frac{dy}{dx} \right)$$

$$\frac{dQ}{dx} = x^y \left[\frac{y}{x} + \log x \frac{dy}{dx} \right] \quad - (2)$$

$$Q = y^x$$

$$R = y^x$$

$$\log R = x \log y$$

$$\frac{1}{R} \frac{dR}{dx} = \frac{x}{y} \frac{dy}{dx} + \log y$$

$$\frac{dx}{dy} = y^x \left[\frac{x}{y} \frac{dy}{dx} + \log y \right] \quad \text{--- (3)}$$

adding ① & ② & ③ we get

$$x^x + x^x \log x + x^{y-1} y + x^y \log x \frac{dy}{dx} + y^{x-1} x \frac{dy}{dx} + y^x \log y = 0$$

$$\frac{dy}{dx} = - \left[x^x (1 + \log x) + x^{y-1} y + y^x \log y \right] \\ x^y \log x + y^{x-1} x$$

Q:18 $y = e^{ax} \cosh bx$

$$\frac{dy}{dx} = e^{ax} (-\sinh bx) b + \cosh bx \times e^{ax} \times a$$

$$\frac{dy}{dx} = -b e^{ax} \sinh bx + a y \Rightarrow \left[\frac{-1}{b} \left(\frac{dy}{dx} - ay \right) = e^{ax} \sinh bx \right] \quad \text{--- (1)}$$

$$\frac{d^2 y}{dx^2} = -b \left[e^{ax} \cosh bx \times b + \sinh bx \times e^{ax} \times a \right] + a \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} = -b \left[by - \frac{a}{b} \left(\frac{dy}{dx} - ay \right) \right] + a \frac{dy}{dx} \quad \text{from ①}$$

$$\frac{d^2y}{dx^2} = -b^2y + a \frac{dy}{dx} - a^2y + a \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} + (a^2 + b^2)y - 2a \frac{dy}{dx} = 0$$

hence proved.

$$\underline{\underline{26^o}} \quad Z = 5x + 2y$$

↓
objective function to be made maximum and minimum

$$x - 2y \leq 2$$

$$3x + 2y \leq 12$$

$$-3x + 2y \leq 3$$

$$x \geq 0, y \geq 0$$

corresponding equations

$$x - 2y = 2$$

put $(0, 0)$ in it

$$0 - 0 \leq 2$$

which is true so the region is towards

origin

$$3x + 2y \leq 12 \Rightarrow 3x + 2y = 12$$

put $(0, 0)$

$$0 \leq 12 \text{ which is true}$$

so region towards origin

$$-3x + 2y = 3$$

put $(0, 0)$

$$0 \leq 3$$

which is true

so region is towards origin

$x \geq 0$ $y \geq 0$. so it means I st quadrant.

at A (2,

at B

at C

at

intersection point of intersection point

$$-3x + 2y = 3$$

$$3x + 2y = 12$$

$$4y = 15$$

$$y = \frac{15}{4}$$

$$-3x + 2 \times \frac{15}{4} = 3$$

$$-3x = 3 - \frac{15}{2}$$

$$-x = 1 - \frac{5}{2}$$

$$-x = -\frac{3}{2}$$

$$x = \frac{3}{2}$$

$$Z = 5x + 2y$$

$$\text{at } A(2, 0) \quad Z = 10$$

$$\text{at } B(0, 0) \quad Z = 0$$

$$\text{at } C(0, \frac{3}{2}) \quad Z = 3$$

$$\text{at } D(\frac{3}{2}, \frac{15}{4}) \quad Z = \frac{15}{2} + \frac{15}{2} = 15$$

of:

$$x - 2y = 2$$

$$3x + 2y = 12$$

$$4x = 14$$

$$x = \frac{7}{2}$$

$$\frac{7}{2} - 2y = 2$$

$$\frac{7}{2} - 2 = 2y$$

$$\frac{3}{2} = 2y$$

$$y = \frac{3}{4}$$

at


$$E(\frac{7}{2}, \frac{3}{4})$$

$$Z = \frac{15}{2} + \frac{3}{2}$$

$$Z = 9$$

Q. 2.6

0 small boy = 1 month



As Z is maximum
at $D(3, 15)$

$Z = 15$

Z is minimum at $(0, 0)$

$[Z \leq 0]$

170

	Jams	Mats	Toys	
$A =$	30	12	70	$\rightarrow$ School X
	40	15	55	$\rightarrow$ School Y
	35	20	75	$\rightarrow$ School Z

3×3

$B =$

25	$\rightarrow$ cost of Jams
100	$\rightarrow$ cost of Mats
50	$\rightarrow$ cost of Toys

cost of Jams

cost of Mats

cost of Toys

$$AB = \begin{bmatrix} 30 & 12 & 70 \\ 40 & 15 & 55 \\ 30 & 20 & 75 \end{bmatrix} \begin{bmatrix} 25 \\ 100 \\ 50 \end{bmatrix}$$

$$X = AB = \begin{bmatrix} 750 + 1200 + 3500 \\ 1000 + 1500 + 2750 \\ 875 + 2000 + 3750 \end{bmatrix}$$

$$X = \begin{bmatrix} 5450 \\ 5250 \\ 6625 \end{bmatrix} \begin{array}{l} \rightarrow \text{fund collected by school X} \\ \rightarrow \text{fund collected by school Y} \\ \rightarrow \text{fund collected by school Z} \end{array}$$

$$\begin{array}{l} \text{fund by } X = \text{Rs. } 5450 \\ \text{fund by } Y = \text{Rs. } 5250 \\ \text{fund by } Z = \text{Rs. } 6625 \end{array}$$

$$\text{Total fund} = \text{Rs. } 17325$$

They are helping victims and hence

the value of helping others is generated.

16°

$$I = \int \frac{x+3}{(x+5)^3} e^x dx$$

$$I = \int \frac{x+5}{(x+5)^3} e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

$$I = \int \frac{1}{(x+5)^2} e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

↓
integrating this by parts is taken as 1st function

$$I = \frac{1}{(x+5)^2} e^x - \int \frac{d}{dx} \left(\frac{1}{(x+5)^2} \right) e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

$$I = \frac{1}{(x+5)^2} e^x + \int \frac{2}{(x+5)^3} e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

$$\text{Q. 1} = \frac{e^x}{(x+5)^2} + c$$

15°

$$x = a \sin \omega t (1 + \cos \omega t)$$

$$\frac{dx}{dt} = a [\dot{\sin \omega t} (-\sin \omega t) (\omega) + (1 + \cos \omega t) \cos \omega t \times \omega]$$

$$\frac{dx}{dt} = 2a [\cos^2 \omega t + \cos \omega t - \sin^2 \omega t]$$

$$\frac{dx}{dt} = 2a [\cos 4t + \cos \omega t] \quad [\cos^2 x - \sin^2 x = \cos 2x]$$

$$y = b \cos \omega t (1 + \cos \omega t)$$

$$y = b \cos \omega t - b \cos^2 \omega t$$

$$\frac{dy}{dt} = -b \sin \omega t \times \omega + b \times 2 \cos \omega t \sin \omega t \times \omega$$

$$\frac{dy}{dt} = 2b [-\sin \omega t + \sin 4t]$$

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$$

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$$

$$\frac{dy}{dx} = \frac{2b}{2a} \left[\frac{\sin 4t - \sin 2t}{\cos 2t + \cos 4t} \right]$$

$$\text{at } t = \frac{\pi}{4}$$

$$\frac{dy}{dx} = \frac{b}{a} \left[\frac{\sin \pi - \sin \frac{\pi}{2}}{\cos \frac{\pi}{2} + \cos \pi} \right]$$

$$= \frac{b}{a} \left[\frac{0 - 1}{0 - 1} \right]$$

$$\frac{dy}{dx} \bigg|_{t=\frac{\pi}{4}} = \frac{b}{a}$$

140

$$I = \frac{1}{2} \int_0^{\frac{\pi}{2}} \frac{\sin^2 x}{\sin x + \cos x} dx \quad \text{--- (1)}$$

$$\left[\int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$I = \frac{1}{2} \int_0^{\frac{\pi}{2}} \frac{\sin^2(\frac{\pi}{2} - x)}{\sin(\frac{\pi}{2} - x) + \cos(\frac{\pi}{2} - x)} dx$$

$$I = \frac{1}{2} \int_0^{\frac{\pi}{2}} \frac{\cos^2 x}{\cos x + \sin x} dx \quad \text{--- (2)}$$

adding (1) & (2) we get

$$2I = \frac{1}{2} \int_0^{\frac{\pi}{2}} \frac{\sin^2 x + \cos^2 x}{\sin x + \cos x} dx$$

$$2I = \frac{1}{2} \int_0^{\frac{\pi}{2}} \frac{1}{(\sin x + \cos x)} dx$$

$$2I = \int_0^1 \frac{1 + \tan^2 x}{2 \tan x + 1 - \tan^2 x} dx$$

$$2I = \int_0^1 \frac{\sec^2 x}{2 \tan x + 1 - \tan^2 x} dx \quad \left\{ \begin{array}{l} \text{put } \sin = 2 \tan x \\ \text{and } \cos x = \frac{1 + \tan^2 x}{1 - \tan^2 x} \end{array} \right.$$

put $\tan x = t$

$$\sec^2 x \times \frac{1}{2} dx = dt$$

$$\begin{array}{l} x \rightarrow 0 \quad t \rightarrow 0 \\ \text{when } x \rightarrow \frac{\pi}{4} \quad t \rightarrow 1 \end{array}$$

$$2I = \int_0^1 \frac{dt}{2t + 1 - t^2}$$

$$I = \int_0^1 \frac{dt}{2t + 1 - t^2} \Rightarrow I = \int_0^1 \frac{dt}{2(t-1)^2}$$

$$I = \int_0^1 \frac{dx}{(\sqrt{x})^2 - (x-1)^2} = \int_0^1 \frac{1}{a^2 - x^2} = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{x} + x - 1}{\sqrt{x} - x + 1} \right| \Bigg|_0^1$$

$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} + 1 - 1}{\sqrt{2} - 1 + 1} \right| - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log 1 - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

$$= 0 - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

$$I = -\frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

13°

$$\Delta = \begin{vmatrix} x+2 & x+6 & x-1 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$R_1 \rightarrow R_1 + R_2 + R_3$$

$$\Delta = \begin{vmatrix} 3x+7 & 3x+7 & 3x+7 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$\Delta = 3x+7 \begin{vmatrix} 1 & 1 & 1 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$C_1 \rightarrow C_1 \rightarrow C_2 \quad C_2 \rightarrow C_2 \rightarrow C_3$$

$$\Delta = (3x+7) \begin{vmatrix} 0 & 0 & 1 \\ 7 & -3 & x+2 \\ -3 & -4 & x+6 \end{vmatrix}$$

expanding by R_1

$$\Delta = (3x+7) [-28 \times 9]$$

$$\Delta = 0$$

$$\Rightarrow 3x+7=0$$

$$x = -\frac{7}{3}$$

120

$$A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$A^2 = A \cdot A =$$

$$\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1+4+4 & 2+2+4 & 2+4+2 \\ 2+2+4 & 4+1+4 & 4+2+2 \\ 2+4+2 & 4+2+2 & 4+4+1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 8 \\ 8 & 8 & 9 \end{bmatrix}$$

$$A^2 - 4A - 5I = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 9 \\ 8 & 8 & 9 \end{bmatrix} + \begin{bmatrix} -4 & -8 & -8 \\ -8 & -4 & -8 \\ -8 & -8 & -4 \end{bmatrix} + \begin{bmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 0 & -5 \end{bmatrix}$$

$$A^2 - 4A - 5I = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = 0$$

$$\text{Hence } A^2 - 4A - 5I = 0$$

Hence proved.

$$A^2 - 4A - 5I = 0$$

pre-multiplying by A^{-1}

$$A^{-1}AA - 4A^{-1}A - 5A^{-1}I = 0$$

$$A^{-1}A = I$$

$$IA - 4I - 5A^{-1} = 0$$

$$IA = A$$

$$A - 4I - 5A^{-1} = 0$$

$$A - 4I = 5A^{-1}$$

$$A^{-1} = \frac{A - 4I}{5}$$

5

$$A^{-1} = \frac{1}{5} [A - 4I]$$

$$5A^{-1} =$$

$$\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} + \begin{bmatrix} -4 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & -4 \end{bmatrix}$$

$$5A^{-1} =$$

$$\begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

$$\text{So } A^{-1} = \frac{1}{5} \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

11.

$$\sin^{-1}(1-x) - 2\sin^{-1}x = \frac{\pi}{2}$$

$$\sin^{-1}(1-x) = \frac{\pi}{2} + 2\sin^{-1}x$$

$$(1-x) = \sin\left(\frac{\pi}{2} + 2\sin^{-1}x\right)$$

$$(1-x) = \cos(2\sin^{-1}x)$$

$$2\sin^{-1}x = 0$$

$$x = \sin 0$$

$$\cos 2\theta = 1 - 2\sin^2\theta$$

$$\cos 2\theta = 1 - 2x^2$$

$\Rightarrow$

$$1-x = \cos 2\theta$$

$$1-x = 1-2x^2$$

$$\Rightarrow 2x^2 - x = 0$$

$$x[2x-1] = 0$$

$$x=0 \text{ or } x=\frac{1}{2}$$

put $x=\frac{1}{2}$ in equation

$$x \sin^{-1} \frac{1}{2} - 2x \sin^{-1} \frac{1}{2}$$

$$= \frac{1}{6} - 2 \times \frac{1}{6}$$

$$\neq \frac{1}{2}$$

$$\text{So } x \neq \frac{1}{2}$$

$$\text{So } \boxed{x=0}$$

passing point of line = $(492, 2)$
since it is // to line

so direction ratio of line $\propto \langle 2, 13, 6 \rangle$

10.

the equation of line

$$\frac{x-4}{2} = \frac{y-2}{3} = \frac{z-2}{6} = \lambda$$

general point on line

$$(2\lambda+4, 3\lambda+2, 6\lambda+2)$$

$$P(1, 2, 3)$$

Q

line

Q be the foot of LP

PQ is $\perp$ to line

the dot product of direction ratios of line and PQ is 0

$$\text{direction } Q = \langle 2\lambda+4, 3\lambda+2, 6\lambda+2 \rangle$$

ratio of

As according to question:

$$2(2\lambda + 3) + (3\lambda)3 + 6(6\lambda - 1) = 0$$

$$4\lambda + 6 + 9\lambda + 36\lambda - 6 = 0$$

$$\boxed{\lambda = 0}$$

As point $Q = (4, 2, 2)$

As for distance

$$PQ = \sqrt{(4-1)^2 + (2-2)^2 + (2-3)^2}$$

$$= \sqrt{(3)^2 + (0)^2 + (-1)^2}$$

$$= \sqrt{9+1}$$

$$\boxed{\text{Length of } PQ = \sqrt{10} \text{ unit}}$$

Qo

~~AB~~

then A, B, C, D are coplanar.

So

$\vec{AB} \cdot (\vec{BC} \times \vec{CD}) = 0$
triple product is 0.

$$\vec{AB} = 1\hat{j} + (x-1)\hat{j} + 4\hat{k}$$

$$\vec{BC} = 0\hat{j} + (1-x)\hat{j} - 7\hat{k}$$

$$\vec{CD} = 2\hat{j} + 3\hat{j} + \hat{k}$$

$$\vec{AB} \cdot (\vec{BC} \times \vec{CD}) = \begin{vmatrix} 1 & x-1 & 4 \\ 0 & 1-x & -7 \\ 2 & 3 & 1 \end{vmatrix} = 0$$

$$\begin{vmatrix} 1 & x-1 & 4 \\ 0 & 1-x & -7 \\ 2 & 3 & 1 \end{vmatrix}$$

expanding by R_1

$$1(1-x+21) - (x-1)14 + 4(2(x-1)) = 0$$

$$22-x - 14x + 14 + 8x - 8 = 0$$

$$-7x = -28 \quad \boxed{x=4}$$

8°

$P(\text{probability of success}) = \frac{1}{2}$
 i.e. that head comes

$q(\text{probability of failure}) = \frac{1}{2}$
 i.e. that tail comes

Let the coin be tossed n times

this event follows the conditions of Bernoulli trial

X be the random variable = no. of heads

$$P(X \geq 1) = P(1) + \dots + P(n)$$

$$P(X \geq 1) = 1 - P(0) \\ = 1 - {}^nC_0 \times \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^n$$

$P(X \geq 8)$ should be more than 80%.

$$P(X \geq 1) > \frac{8}{10} \\ \frac{86}{100} < 1 - \left(\frac{1}{2}\right)^n$$

$$1 - \left(\frac{1}{2}\right)^n > \frac{8}{10}$$

$$\left(1 - \frac{8}{10}\right)^n > \left(\frac{1}{2}\right)^n$$

$$\frac{18}{5} > \left(\frac{1}{2}\right)^n$$

$$5 < 2^n$$

$$n = 3$$

follow the condition
So the coin should be tossed
at least 3 times.

Ex.

$$I = \int \frac{x^2}{x^4 + x^2 - 2} dx$$

$$I = \int \frac{x^2}{x^4 + 2x^2 - x^2 - 2} dx$$

$$I = \int \frac{x^2}{x^2(x^2 + 2) - 1(x^2 + 2)} dx$$

$$I = \int \frac{x^2}{(x^2-1)(x^2+2)} dx$$

put $x^2 = t$

then $\frac{t}{(t-1)(t+2)} = \frac{A}{t-1} + \frac{B}{t+2}$

$$t = A(t+2) + B(t-1)$$

put $t=1$

$$1 = A \times 3$$

$$\boxed{A = \frac{1}{3}}$$

put $t=-2$

$$-2 = -3B$$

$$\boxed{B = \frac{2}{3}}$$

$$I = \int \frac{1}{3(t-1)} dt + \frac{2}{3} \int \frac{dx}{t+2}$$

$$I = \int \frac{1}{3(x^2-1)} dx + \frac{2}{3} \int \frac{dx}{(x^2+2)} \quad \int \frac{1}{x^2-a^2} = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right|$$

$$I = \frac{1}{3} \times \frac{1}{2} \log \left| \frac{x-1}{x+1} \right| + \frac{2}{3} \times \frac{1}{\sqrt{2}} \tan^{-1} \frac{x}{\sqrt{2}} + C \quad \int \frac{1}{x^2+a^2} = \frac{1}{a} \tan^{-1} \frac{x}{a}$$

$$I = \frac{1}{6} \log \left| \frac{x-1}{x+1} \right| + \frac{\sqrt{2}}{3} \tan^{-1} \frac{x}{\sqrt{2}} + C$$

Section - A

Q: 6

equation of plane

$$6x - 3y + 2z - 4 = 0$$

distance =

$$\frac{|6 \times 2 - 3 \times 5 + 2(-3) - 4|}{\sqrt{36 + 9 + 4}}$$

$$\frac{|12 - 15 - 6 - 4|}{\sqrt{49}}$$

$$= \frac{|12 - 15 - 10|}{7}$$

$$\frac{\text{distance}}{7} = \frac{13}{7} \text{ units}$$

Q: 5

$$\vec{a} = \hat{i} - \hat{j}$$

$$|\vec{a}| = \sqrt{2}$$

$$\vec{b} = \hat{j} - \hat{k}$$

$$|\vec{b}| = \sqrt{2}$$

$$|\vec{a} \times \vec{b}| = \sqrt{(-17)^2 + (13)^2 + (7)^2}$$

$$= \sqrt{289 + 169 + 49}$$

$$= \sqrt{507}$$

$$= \sqrt{3 \times 169}$$

$$|\vec{a} \times \vec{b}| = 13\sqrt{3}$$

Q. 3.

$$x \log x \frac{dy}{dx} + y = 2 \log x$$

$$\frac{dy}{dx} + \frac{y}{x \log x} = 2 \log x$$

compare it with $\frac{dy}{dx} + P_y = Q$

$$I.F. =$$

$$e^{\int P dx}$$

$$= e^{\int \frac{1}{x \log x} dx}$$

$$\text{put } \log x = t$$

$$\frac{1}{x} dx = dt$$

$$I \circ F = e^{\int \frac{dt}{t}}$$

$$= e^{\log |t|}$$

$$= t$$

$$I \circ F = \log x$$

2°

general equation of family of lines passing through origin

$$y = mx$$

$$m = \frac{y}{x}$$

$$\frac{dy}{dx} = m$$

$$\frac{dy}{dx} = \frac{y}{x}$$

$$x \frac{dy}{dx} - y = 0$$

Q11

$$A_{12} = e^{2 \times 1 \times 2} \sin 2x$$

$$A_{12} = e^{2x} \sin 2x$$

~~Q11~~

~~Question~~

~~done as per~~

~~01/07/2022~~

~~Marked~~

XIV - U